

_		Allergies:		
Date of Birth:				
Surgeon:				
Date of Procedure:				
Procedure:				
1				
PRE-ADMISSION TESTING ORDERS				
l				
LABS AND TESTIN		RADIOLOGY:	cay (CVP)	
	BMP LABS/TESTING PER ANESTHESIA GUIDELINES CBC	2 View chest x-r	ay (CXR)	
	CMP	PER STANDING DELEGAT	TED ORDER:	
	HCT	12 LEAD EKG		
	HGB	MRSA SCREENIN	IG	
	HbA1c	PREGNANCY TE	EST URINE SERUM	
	Prothrombin Time (PT)	TYPE and SCRE	EN	
	Partial Thromboplastin Time (PTT)	TYPE and RH		
	SED Rate			
	MRSA PCR			
	Other TAH major GVN and Colorectal current TAH major GVN and Colorectal cu	raorios:		
Use Surgical site Infection (SSI) orders for General surgery, TAH, major GYN and Colorectal surgeries: Draw Liver Panel, CEA, BNP, add HbA1c for TAH, major GYN, and Colorectal surgery				
	Follow ERAS RCB Patient Education and Pre-Operative Instructions for Spine, Total Joint Hip/Knee, and			
	Colorectal Surgery			
	PRE-OPERATIVE ORDERS (Day of Surgery Orders)			
ADMIT:				
	Admit to Inpatient Place in Outpatient for Same Day Procedure/Surgery	☐ Place in Outpatient	WITH overnight Stay	
ANTIBIOTICS:				
	Cefazolin 2gm IVP if <120kg x 1, 3gm if > = 120kg x 1 to be given by Anesthesia			
	Clindamycinmg IVPB x 1 to be given by Anesthesia			
	Gentamicinmg IVPB x 1 to be given by Anesthesia			
	Levofloxacin 500mg IVPB x 1 to infuse over 60 minutes, to be given by Anesthesia			
	Vancomycin 1 gm IVPB x 1 to infuse over 90 minutes, START in pre-op			
	Other			
IV FLUID: Start IV				
Lactated Ringers 1000mL atmL/hr or KVO (20mL/hr). May use up to 0.2mL of 1% Lidocaine intradermal for IV start.				
PREOP MEDICATI	ION:	PER STANDING DELEGAT	FD ORDER:	
TREOF MEDICALI	Acetaminophen mg PO x 1	12 Lead EKG	ED ORDER.	
	Gabapentin mg PO x 1	Pregnancy Test		
	Celecoxib mg PO x 1	Type and Scree		
	Other	Type and RH		
	Other	Glucose (point	of care)	
		MRSA Decoloni		
		Glycemic Mana	gement for Colon and Hysterectomy	
MAJOR JOINT REPLACEMENT (for patients who are MRSA positive/or of unknown status and have not LABS:				
completed a 5-da	y treatment prior to surgery) Choose decolonization treatment.	BMP (Rapid Ca	re Test)	
	Mupirocin 2% (Bactroban) to each nostril Povidone-Iodine nasal swab and CHG bath	HGB (Rapid Ca		
	Povidone-louine hasarswab and Cho bath	☐ MRSA PCR	,	
		STAT		
EMBOLISM PREVENTION:				
	TED Hose			
	SCDs to be placed in OR			
	Lovenox 40mg subcutaneous to be given in OR			
1				
		<u>-</u> .		
MD Signature:		Date:	Time:	